

=> d his

(FILE 'HOME' ENTERED AT 12:28:11 ON 06 MAY 2003)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS,  
LIFESCI' ENTERED AT 12:28:34 ON 06 MAY 2003

L1 10975 S ADENYLATE KINASE?  
L2 2615 S HUMAN AND L1  
L3 613747 S MITOCHONDR?  
L4 255 S L2 AND L3  
L5 5929362 S CLON? OR EXPRESS? OR RECOMBINANT  
L6 59 S L4 AND L5  
L7 32 DUP REM L6 (27 DUPLICATES REMOVED)  
L8 8 S "HMAK"  
L9 3 DUP REM L8 (5 DUPLICATES REMOVED)  
E HILLMAN J L/AU  
L10 454 S E3  
L11 1 S L4 AND L10  
E SHAH P/AU  
L12 1409 S E3  
L13 1 S L4 AND L12

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1652MXM

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 3 Jun 03 New e-mail delivery for search results now available  
NEWS 4 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN  
NEWS 5 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)  
now available on STN  
NEWS 6 Aug 26 Sequence searching in REGISTRY enhanced  
NEWS 7 Sep 03 JAPIO has been reloaded and enhanced  
NEWS 8 Sep 16 Experimental properties added to the REGISTRY file  
NEWS 9 Sep 16 CA Section Thesaurus available in CAPLUS and CA  
NEWS 10 Oct 01 CASREACT Enriched with Reactions from 1907 to 1985  
NEWS 11 Oct 24 BEILSTEIN adds new search fields  
NEWS 12 Oct 24 Nutraceuticals International (NUTRACEUT) now available on STN  
NEWS 13 Nov 18 DKILIT has been renamed APOLLIT  
NEWS 14 Nov 25 More calculated properties added to REGISTRY  
NEWS 15 Dec 04 CSA files on STN  
NEWS 16 Dec 17 PCTFULL now covers WP/PCT Applications from 1978 to date  
NEWS 17 Dec 17 TOXCENTER enhanced with additional content  
NEWS 18 Dec 17 Adis Clinical Trials Insight now available on STN  
NEWS 19 Jan 29 Simultaneous left and right truncation added to COMPENDEX,  
ENERGY, INSPEC  
NEWS 20 Feb 13 CANCERLIT is no longer being updated  
NEWS 21 Feb 24 METADEX enhancements  
NEWS 22 Feb 24 PCTGEN now available on STN  
NEWS 23 Feb 24 TEMA now available on STN  
NEWS 24 Feb 26 NTIS now allows simultaneous left and right truncation  
NEWS 25 Feb 26 PCTFULL now contains images  
NEWS 26 Mar 04 SDI PACKAGE for monthly delivery of multifile SDI results  
NEWS 27 Mar 19 APOLLIT offering free connect time in April 2003  
NEWS 28 Mar 20 EVENTLINE will be removed from STN  
NEWS 29 Mar 24 PATDPAFULL now available on STN  
NEWS 30 Mar 24 Additional information for trade-named substances without  
structures available in REGISTRY  
NEWS 31 Apr 11 Display formats in DGENE enhanced  
NEWS 32 Apr 14 MEDLINE Reload  
NEWS 33 Apr 17 Polymer searching in REGISTRY enhanced  
NEWS 34 Apr 21 Indexing from 1947 to 1956 being added to records in CA/CAPLUS  
NEWS 35 Apr 21 New current-awareness alert (SDI) frequency in  
WPIDS/WPINDEX/WPIX  
NEWS 36 Apr 28 RDISCLOSURE now available on STN  
NEWS 37 May 05 Pharmacokinetic information and systematic chemical names  
added to PHAR

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT  
MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),  
AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003

|            |   |
|------------|---|
| NEWS HOURS | STN Operating Hours Plus Help Desk Availability         |
| NEWS INTER | General Internet Information                            |
| NEWS LOGIN | Welcome Banner and News Items                           |
| NEWS PHONE | Direct Dial and Telecommunication Network Access to STN |
| NEWS WWW   | CAS World Wide Web Site (general information)           |

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:28:11 ON 06 MAY 2003

|  |            |         |
|--|------------|---------|
| => file medline embase biosis biotechds scisearch hcaplus ntis lifesci |            |         |
| COST IN U.S. DOLLARS   | SINCE FILE | TOTAL   |
|  | ENTRY      | SESSION |
| FULL ESTIMATED COST  | 0.21       | 0.21    |

FILE 'MEDLINE' ENTERED AT 12:28:34 ON 06 MAY 2003

FILE 'EMBASE' ENTERED AT 12:28:34 ON 06 MAY 2003  
 COPYRIGHT (C) 2003 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:28:34 ON 06 MAY 2003  
 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'BIOTECHDS' ENTERED AT 12:28:34 ON 06 MAY 2003  
 COPYRIGHT (C) 2003 THOMSON DERWENT AND INSTITUTE FOR SCIENTIFIC INFORMATION

FILE 'SCISEARCH' ENTERED AT 12:28:34 ON 06 MAY 2003  
 COPYRIGHT 2003 THOMSON ISI

FILE 'HCAPLUS' ENTERED AT 12:28:34 ON 06 MAY 2003  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'NTIS' ENTERED AT 12:28:34 ON 06 MAY 2003  
 Compiled and distributed by the NTIS, U.S. Department of Commerce.  
 It contains copyrighted material.  
 All rights reserved. (2003)

FILE 'LIFESCI' ENTERED AT 12:28:34 ON 06 MAY 2003  
 COPYRIGHT (C) 2003 Cambridge Scientific Abstracts (CSA)

=> s adenylate kinase?  
 L1 10975 ADENYLATE KINASE?

=> s human and l1  
 L2 2615 HUMAN AND L1

=> s mitochondr?

L3 613747 MITOCHONDR?

=> s 12 and 13

L4 255 L2 AND L3

=> s clon? or express? or recombinant

3 FILES SEARCHED...

L5 5929362 CLON? OR EXPRESS? OR RECOMBINANT

=> s 14 and 15

L6 59 L4 AND L5

=> dup rem 16

PROCESSING COMPLETED FOR L6

L7 32 DUP REM L6 (27 DUPLICATES REMOVED)

=> d 1-32 ibib ab

L7 ANSWER 1 OF 32 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 2002:637880 HCAPLUS

DOCUMENT NUMBER: 137:179893

TITLE: Methods for identifying compounds that inhibit or reduce PTP1B (protein tyrosine phosphatase 1B)

**expression**

INVENTOR(S): Zinker, Bradley A.; Trevillyan, James M.; Jirousek, Michael R.; Rondinone, Christina M.; Cowser, Lex M.; Wyatt, Jacqueline; Monia, Brett P.; Butler, Madeline M.; Waring, Jeffrey French

PATENT ASSIGNEE(S): Abbott Laboratories, USA; Isis Pharmaceuticals, Inc.

SOURCE: PCT Int. Appl., 72 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO.   | KIND | DATE     | APPLICATION NO. | DATE     |
|--|------|----------|-----------------|----------|
| WO 2002064840  | A2   | 20020822 | WO 2002-US4194  | 20020213 |
| W: CA, JP, MX  |      |          |                 |          |
| RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR |      |          |                 |          |

PRIORITY APPLN. INFO.: US 2001-268399P P 20010213

US 2002-74194 A 20020212

AB The present invention relates to methods for identifying compds. that inhibit PTP1B (protein tyrosine phosphatase 1B) mRNA and protein **expression** in insulin resistant obese non-human mammals. The present invention relates to biol. markers for PTP1B inhibition or redn. Specifically, the present invention relates to methods for measuring the downregulation of the p85.alpha. regulatory subunit of phosphatidylinositol-3-kinase and the upregulation of p55.alpha. and/or p50.alpha. isoforms in response to in vivo inhibition or redn. of PTP1B in insulin resistant mammals. Moreover, the present invention relates to an in vivo marker for pharmacodynamic measurements and mechanism of action detns. of small mol. drugs which inhibit or reduce PTP1B activity. Finally, the present invention also provides a method to screen agents for activity that down modulates p85.alpha. and upregulates phosphatidylinositol-3-kinase p85.alpha. isoforms as drugs for the treatment of type 2 diabetes.

L7 ANSWER 2 OF 32 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 2002:937303 HCAPLUS

DOCUMENT NUMBER: 138:20443  
 TITLE: Endocrine disruptor screening using DNA chips of endocrine disruptor-responsive genes  
 INVENTOR(S): Kondo, Akihiro; Takeda, Takeshi; Mizutani, Shigetoshi; Tsujimoto, Yoshimasa; Takashima, Ryokichi; Enoki, Yuki; Kato, Ikunoshin  
 PATENT ASSIGNEE(S): Takara Bio Inc., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 386 pp.  
 CODEN: JKXXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

| PATENT NO.             | KIND | DATE     | APPLICATION NO. | DATE       |
|------------------------|------|----------|-----------------|------------|
| JP 2002355079          | A2   | 20021210 | JP 2002-69354   | 20020313   |
| PRIORITY APPLN. INFO.: |      |          | JP 2001-73183   | A 20010314 |
|                        |      |          | JP 2001-74993   | A 20010315 |
|                        |      |          | JP 2001-102519  | A 20010330 |

AB A method and kit for detecting endocrine-disrupting chems. using DNA microarrays are claimed. The method comprises prep. a nucleic acid sample contg. mRNAs or cDNAs originating in cells, tissues, or organisms which have been brought into contact with a sample contg. the endocrine disruptor. The nucleic acid sample is hybridized with DNA microarrays having genes affected by the endocrine disruptor or DNA fragments originating in these genes have been fixed. The results obtained are then compared with the results obtained with the control sample to select the gene affected by the endocrine disruptor. Genes whose **expression** is altered by tri-Bu tin, 4-octaphenol, 4-nonylphenol, di-N-Bu phthalate, dichlorohexyl phthalate, octachlorostyrene, benzophenone, diethylhexyl phthalate, diethylstilbestrol (DES), and 17-.beta. estradiol (E2), were found in mice by DNA chip anal.

L7 ANSWER 3 OF 32 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.

ACCESSION NUMBER: 2002449003 EMBASE  
 TITLE: Old and new determinants in the regulation of energy expenditure.  
 AUTHOR: Russell A.P.; Giacobino J.P.  
 CORPORATE SOURCE: Prof. J.P. Giacobino, Departement de Biochimie Medicale, Centre Medical Universitaire, 1 rue Michel-Servet, 1211 Geneve 4, Switzerland. Jean-Paul.Giacobino@medecine.unige.ch  
 SOURCE: Journal of Endocrinological Investigation, (2002) 25/10 (862-866).  
 Refs: 55  
 ISSN: 0391-4097 CODEN: JEIND7  
 COUNTRY: Italy  
 DOCUMENT TYPE: Journal; General Review  
 FILE SEGMENT: 003 Endocrinology  
 029 Clinical Biochemistry  
 005 General Pathology and Pathological Anatomy  
 030 Pharmacology  
 037 Drug Literature Index  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB Bw gain is controlled by energy intake on one hand and expenditure on the other. The components of energy expenditure are basal metabolism, exercise induced thermogenesis and adaptive thermogenesis. In this short review we shall discuss the main determinants of adaptive thermogenesis.  
 .COPYRG.T.2002, Editrice Kurtis.

L7 ANSWER 4 OF 32 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.  
ACCESSION NUMBER: 2002446521 EMBASE  
TITLE: Frontiers in research on parasitic protozoa.  
AUTHOR: Gibson W.; Miles M.  
CORPORATE SOURCE: W. Gibson, School of Biological Sciences, University of  
Bristol, Woodland Road, Bristol BS8 1UG, United Kingdom.  
w.gibson@bristol.ac.uk  
SOURCE: Trends in Parasitology, (1 Dec 2002) 18/12 (521-522).  
Refs: 2  
ISSN: 1471-4922 CODEN: TPRACT  
PUBLISHER IDENT.: S 1471-4922(02)02416-9  
COUNTRY: United Kingdom  
DOCUMENT TYPE: Journal; Conference Article  
FILE SEGMENT: 004 Microbiology  
037 Drug Literature Index  
LANGUAGE: English

L7 ANSWER 5 OF 32 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
ACCESSION NUMBER: 2002-00258 BIOTECHDS  
TITLE: New antibody against **human mitochondria**  
**adenylate-kinase** isozyme 2 or isozyme 3,  
for detecting the isozymes in a detection sample to diagnose  
cardiac diseases such as myocardial infarction and angina  
pectoris;  
monoclonal antibody, hybridoma cell culture and detection  
marker useful in disease diagnosis  
AUTHOR: Cho K S; Lee S M  
PATENT ASSIGNEE: Kim H J  
LOCATION: Ansan, Korea.  
PATENT INFO: WO 2001058482 16 Aug 2001  
APPLICATION INFO: WO 2000-KR882 10 Aug 2000  
PRIORITY INFO: KR 2000-5808 8 Feb 2000  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
OTHER SOURCE: WPI: 2001-522438 [57]

AB An antibody (I) specific to **human mitochondria**  
**adenylate-kinase** (AK) isozymes AK2 or AK3 or their  
portion, is claimed. (I) is produced in an animal species and has a  
reactivity with the immunogen which includes a **human**  
**mitochondria adenylate-kinase** isozyme or its  
portion. Also claimed are: an immunological formulation (II) for  
diagnosing cardiac disease containing (I) and a detection marker; and a  
diagnostic kit (III) for cardiac disease containing a carrier and (I)  
which is coupled with a detection marker. (I) is useful for detecting a  
**human mitochondrial adenylate-kinase**  
isozyme (AK2) or (AK3) in a detection sample. An immunological  
formulation (II) for diagnosing cardiac disease containing (I) and a  
detection marker is useful for detecting **adenylate-**  
**kinase** isozyme in a biological sample. (I) is useful for  
diagnosing cardiac disease such as myocardial infarction, angina  
pectoris. (II) and a diagnostic kit (III) are also useful for  
diagnosing cardiac disease. (56pp)

L7 ANSWER 6 OF 32 HCAPLUS COPYRIGHT 2003 ACS  
ACCESSION NUMBER: 2001:597830 HCAPLUS  
DOCUMENT NUMBER: 135:194482  
TITLE: Anti-**human mitochondrial**  
**adenylate kinase** isozyme antibody,  
diagnostic formulation and diagnostic kit for cardiac  
disease  
INVENTOR(S): Kim, Hyo Joon; Cho, Key Seung; Lee, Sang Min  
PATENT ASSIGNEE(S): S. Korea

SOURCE: PCT Int. Appl., 56 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

| PATENT NO.   | KIND | DATE     | APPLICATION NO. | DATE       |
|--|------|----------|-----------------|------------|
| WO 2001058482  | A1   | 20010816 | WO 2000-KR882   | 20000810   |
| W: AU, BR, CA, CN, DE, JP, RU, US  |      |          |                 |            |
| RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE |      |          |                 |            |
| EP 1165133   | A1   | 20020102 | EP 2000-955110  | 20000810   |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI  |      |          |                 |            |
| BR 2000010712  | A    | 20020205 | BR 2000-10712   | 20000810   |
| PRIORITY APPLN. INFO.:   |      |          | KR 2000-5808    | A 20000208 |
|  |      |          | WO 2000-KR882   | W 20000810 |

AB The present invention relates to an immunol. formulation and a diagnostic kit for cardiac disease using **human mitochondrial adenylate kinase** isoenzymes. The present invention provides an immunol. formulation and a diagnostic kit for cardiac disease, which are featured by using **mitochondrial adenylate kinase** isoenzymes which exist in a myocardial cell among muscle cells, but not in a skeletal muscular cell, as a diagnostic marker for cardiac disease and which enable more correct and easy diagnosis of cardiac disease.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 7 OF 32 MEDLINE DUPLICATE 1  
 ACCESSION NUMBER: 2001439872 MEDLINE  
 DOCUMENT NUMBER: 21378190 PubMed ID: 11485571  
 TITLE: Structure and **expression** of **human mitochondrial adenylate kinase** targeted to the **mitochondrial** matrix.  
 AUTHOR: Noma T; Fujisawa K; Yamashiro Y; Shinohara M; Nakazawa A; Gondo T; Ishihara T; Yoshinobu K  
 CORPORATE SOURCE: Department of Biochemistry, Yamaguchi University School of Medicine, 1-1-1 Minami-Kogushi, Ube, Yamaguchi 755-8505, Japan.. tnoma@po.cc.yamaguchi-u.ac.jp  
 SOURCE: BIOCHEMICAL JOURNAL, (2001 Aug 15) 358 (Pt 1) 225-32.  
 Journal code: 2984726R. ISSN: 0264-6021.  
 PUB. COUNTRY: England: United Kingdom  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-AB021870  
 ENTRY MONTH: 200109  
 ENTRY DATE: Entered STN: 20010924  
 Last Updated on STN: 20010924  
 Entered Medline: 20010920

AB The previously isolated cDNA encoding **human adenylate kinase** (AK) isozyme 3 was recently renamed AK4. Consequently, **human** AK3 cDNA remains to be identified and we have little information about the functional relationship between **human** AK3 and AK4. In pursuit of the physiological roles of both the AK3 and AK4 proteins, we first isolated an authentic **human** AK3 cDNA and compared their **expression**. Nucleotide sequencing revealed that the cDNA encoded a 227-amino-acid protein, with a deduced molecular mass of 25.6 kDa, that shares greater homology with the AK3 cDNAs isolated from

bovine and rat than that from **human**. We named the isolated cDNA AK3. Northern-blot analysis revealed that AK3 mRNA was present in all tissues examined, and was highly **expressed** in heart, skeletal muscle and liver, moderately **expressed** in pancreas and kidney, and weakly **expressed** in placenta, brain and lung. On the other hand, we found that **human** AK4 mRNA was highly **expressed** in kidney, moderately **expressed** in heart and liver and weakly **expressed** in brain. Western-blot analysis demonstrated **expression** profiles of AK3 and AK4 that were similar to their mRNA **expression** patterns in each tissue. Over **expression** of AK3, but not AK4, in both Escherichia coli CV2, a temperature-sensitive AK mutant, and a **human** embryonic kidney-derived cell line, HEK-293, not only produced significant GTP:AMP phosphotransferase (AK3) activity, but also complemented the CV2 cells at 42 degrees C. Subcellular and submitochondrial fractionation analysis demonstrated that both AK3 and AK4 are localized in the **mitochondrial** matrix.

L7 ANSWER 8 OF 32 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 ACCESSION NUMBER: 2002:288316 BIOSIS  
 DOCUMENT NUMBER: PREV200200288316  
 TITLE: Hemodynamic unloading by ventricular assist devices has no beneficial effect for the inflammation-associated apoptotic pathway in **human** terminally failing myocardium.  
 AUTHOR(S): Scheubel, Robert Johannes (1); Bartling, Babett (1); Stein, Susanne; Darmer, Dorothea; Holtz, Juergen; Silber, Rolf-Edgar  
 CORPORATE SOURCE: (1) Clin fuer Herz- und Thoraxchirurgie, Halle/Saale Germany  
 SOURCE: Circulation, (October 23, 2001) Vol. 104, No. 17 Supplement, pp. II.713. <http://circ.ahajournals.org/>. print.  
 Meeting Info.: Scientific Sessions 2001 of the American Heart Association Anaheim, California, USA November 11-14, 2001  
 ISSN: 0009-7322.  
 DOCUMENT TYPE: Conference  
 LANGUAGE: English

L7 ANSWER 9 OF 32 HCAPLUS COPYRIGHT 2003 ACS  
 ACCESSION NUMBER: 2000:810680 HCAPLUS  
 DOCUMENT NUMBER: 133:345587  
 TITLE: Protein and cDNA sequences of a novel **human Mitochondria adenylate kinase** GTP3P and uses thereof  
 INVENTOR(S): Yu, Long; Zhao, Yong; Bi, Anding; Gao, Jie; Zhao, Shouyuan  
 PATENT ASSIGNEE(S): Fudan Gene Engineering Co., Ltd., Xinhuangpu, Shanghai, Peop. Rep. China  
 SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 20 pp. CODEN: CNXXEV  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Chinese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| CN 1249340  | A    | 20000405 | CN 1998-119439  | 19980928 |
| PRIORITY APPLN. INFO.:  |      |          | CN 1998-119439  | 19980928 |
| AB The invention provides protein and cDNA sequences of a novel <b>human Mitochondria adenylate kinase</b> GTP3P which is belived to be a GTP-AMP transphosphorylase. The invention also relates to |      |          |                 |          |



constructing **adenylate kinase** GTP3P expression cassette to producing **recombinant adenylate kinase** GTP3P using E.coli cells or eukaryotic cells. The invention further relates to the uses of **adenylate kinase** GTP3P.

L7 ANSWER 10 OF 32 SCISEARCH COPYRIGHT 2003 THOMSON ISI  
ACCESSION NUMBER: 2000:361124 SCISEARCH  
THE GENUINE ARTICLE: 311BC  
TITLE: Cellular phosphorylation of 2',3'-dideoxyadenosine-5'-monophosphate, a key intermediate in the activation of the antiviral agent DDI, in human peripheral blood mononuclear cells  
AUTHOR: Robbins B L (Reprint); Greenshaw J; Fridland A  
CORPORATE SOURCE: ST JUDE CHILDRENS HOSP, DEPT INFECT DIS, 332 N LAUDERDALE ST, MEMPHIS, TN 38105 (Reprint)  
COUNTRY OF AUTHOR: USA  
SOURCE: NUCLEOSIDES NUCLEOTIDES & NUCLEIC ACIDS, (MAY 2000) Vol. 19, No. 1-2, pp. 405-413.  
Publisher: MARCEL DEKKER INC, 270 MADISON AVE, NEW YORK, NY 10016.  
ISSN: 1525-7770.  
DOCUMENT TYPE: Article; Journal  
FILE SEGMENT: LIFE  
LANGUAGE: English  
REFERENCE COUNT: 20

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB 2',3'-dideoxyadenosine 5-monophosphate (ddAMP), is a key intermediate in the metabolism of the antiviral agent 2',3'-dideoxyinosine (ddI) to its active triphosphate derivative, 2',3'-dideoxyadenosine-5'-triphosphate (ddATP). The potential role of **adenylate kinase** in the phosphorylation of ddAMP was studied in human peripheral blood mononuclear cells (PBMC) and a human T cell line, CEMss. Subcellular distribution, sulfhydryl inhibitor, and substrate specificity studies support the hypothesis that the **mitochondrial adenylate kinase** (AK2) is a major route of cellular activation of these compounds in human lymphocytes.

L7 ANSWER 11 OF 32 MEDLINE DUPLICATE 2  
ACCESSION NUMBER: 2000246295 MEDLINE  
DOCUMENT NUMBER: 20246295 PubMed ID: 10786623  
TITLE: cDNA **cloning** and chromosomal mapping of the gene encoding **adenylate kinase** 2 from Drosophila melanogaster.  
AUTHOR: Noma T; Murakami R; Yamashiro Y; Fujisawa K; Inouye S; Nakazawa A  
CORPORATE SOURCE: Department of Biochemistry, Yamaguchi University School of Medicine, Ube, Japan.. tnoma@po.cc.yamaguchi-u.ac.jp  
SOURCE: BIOCHIMICA ET BIOPHYSICA ACTA, (2000 Jan 31) 1490 (1-2) 109-14.  
Journal code: 0217513. ISSN: 0006-3002.  
PUB. COUNTRY: Netherlands  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
OTHER SOURCE: GENBANK-AB009996; GENBANK-AC004642  
ENTRY MONTH: 200005  
ENTRY DATE: Entered STN: 20000606  
Last Updated on STN: 20000606  
Entered Medline: 20000524

AB As a step toward understanding of the role of **adenylate kinase** (AK) in energy metabolism, we analyzed this enzyme in

*Drosophila melanogaster*. The enzyme activities of all three AK isozymes were determined in cell-free extracts of flies, and their proteins were detected by Western blot analysis using polyclonal antibodies against the mammalian isozymes. A cDNA encoding **adenylate kinase** was isolated from *D. melanogaster* cDNA library. The cDNA encodes a 240-amino acid protein, which shows high similarity to bovine, **human** and rat AK2, and hence was named DAK2. Preliminary subcellular fractionation analysis indicated that DAK2 is localized in both cytoplasm and **mitochondria**. In situ hybridization to salivary gland polytene chromosomes revealed that the *Dak2* gene is located at 60B on the right arm of the second chromosome.

L7 ANSWER 12 OF 32 SCISEARCH COPYRIGHT 2003 THOMSON ISI  
 ACCESSION NUMBER: 2000:126507 SCISEARCH  
 THE GENUINE ARTICLE: 282KY  
 TITLE: cDNA **cloning** and chromosomal mapping of the gene encoding **adenylate kinase** 2 from *Drosophila melanogaster*  
 AUTHOR: Noma T (Reprint); Murakami R; Yamashiro Y; Fujisawa K; Inouye S; Nakazawa A  
 CORPORATE SOURCE: YAMAGUCHI UNIV, SCH MED, DEPT BIOCHEM, YAMAGUCHI 7558505, JAPAN (Reprint); YAMAGUCHI UNIV, FAC SCI, DEPT PHYS BIOL & INFORMAT, YAMAGUCHI 7538512, JAPAN  
 COUNTRY OF AUTHOR: JAPAN  
 SOURCE: BIOCHIMICA ET BIOPHYSICA ACTA-GENE STRUCTURE AND EXPRESSION, (31 JAN 2000) Vol. 1490, No. 1-2, pp. 109-114. Publisher: ELSEVIER SCIENCE BV, PO BOX 211, 1000 AE AMSTERDAM, NETHERLANDS. ISSN: 0167-4781.  
 DOCUMENT TYPE: Article; Journal  
 FILE SEGMENT: LIFE  
 LANGUAGE: English  
 REFERENCE COUNT: 32

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB As a step toward understanding of the role of **adenylate kinase** (AK) in energy metabolism, we analyzed this enzyme in *Drosophila melanogaster*. The enzyme activities of all three AK isozymes were determined in cell-free extracts of flies, and their proteins were detected by Western blot analysis using polyclonal antibodies against the mammalian isozymes. A cDNA encoding **adenylate kinase** was isolated from *D. melanogaster* cDNA library. The cDNA encodes a 240-amino acid protein, which shows high similarity to bovine, **human** and rat AK2, and hence was named DAK2. Preliminary subcellular fractionation analysis indicated that DAK2 is localized in both cytoplasm and **mitochondria**. In situ hybridization to salivary gland polytene chromosomes revealed that the *Dak2* gene is located at 60B on the right arm of the second chromosome. (C) 2000 Elsevier Science B.V. All rights reserved.

L7 ANSWER 13 OF 32 HCAPLUS COPYRIGHT 2003 ACS  
 ACCESSION NUMBER: 2001:782966 HCAPLUS  
 DOCUMENT NUMBER: 136:322434  
 TITLE: **Expression** of mRNAs encoding **adenylate kinase** isozymes 1, 2, 3, and 4 in mouse tissues and during neuronal differentiation of P19 embryonal carcinoma cells  
 AUTHOR(S): Yamashiro, Yasuhiro  
 CORPORATE SOURCE: Department of Biochemistry, Yamaguchi University School of Medicine, Yamaguchi, 755-8505, Japan  
 SOURCE: Bulletin of the Yamaguchi Medical School (2000), 47(3-4), 55-68  
 CODEN: BYMSAN; ISSN: 0513-1812

PUBLISHER: Yamaguchi University, School of Medicine  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The authors **cloned** cDNAs encoding four **adenylate kinase** (AK) isoenzymes from a mouse kidney cDNA library. The AK1, AK2, AK3, and AK4 cDNAs encode the 194-, 232-, 227-, and 223-amino acid proteins, resp. AK4 is a recently isolated gene that is highly homologous to the reported **human** AK3. Northern blot anal. and reverse transcription-polymerase chain reaction anal. revealed that AK1 mRNA was predominantly **expressed** in skeletal muscle, heart, and testis; AK2 mRNA in liver, heart, kidney, and testis; AK3 mRNA almost uniformly in all tissues examd.; and AK4 mRNA prominently in kidney. Subcellular and submitochondrial fractionation anal. suggested that AK4 was localized in the **mitochondrial** matrix. Further, the authors found a 76-fold induction of AK1 mRNA **expression** concomitant with a 53-fold induction of NeuroD **expression** during retinoic acid-induced neuronal differentiation of P19 embryonic carcinoma cell. AK2 and AK3 mRNA **expression** was increased by 4- to 6-fold during differentiation, whereas AK4 transcription was first down-regulated and subsequently returned to the original level. These data on AK isoenzyme gene **expression** may provide basic information for prodn. and evaluation of transgenic mice as well as knockout mice to further understand the **physiol.** role of AK isoenzymes.

REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 14 OF 32 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2000:290852 BIOSIS

DOCUMENT NUMBER: PREV200000290852

TITLE: **Mitochondrial adenylate kinase**

AUTHOR(S): Hillman, Jennifer L.; Shah, Purvi  
ASSIGNEE: Incyte Pharmaceuticals, Inc.

PATENT INFORMATION: US 6001624 December 14, 1999

SOURCE: Official Gazette of the United States Patent and Trademark Office Patents, (Dec. 14, 1999) Vol. 1229, No. 2, pp. No pagination. e-file.  
ISSN: 0098-1133.

DOCUMENT TYPE: Patent

LANGUAGE: English

AB The present invention provides a **human mitochondrial adenylate kinase** (HMAK) and polynucleotides which encode HMAK. The invention also provides **expression** vectors, host cells, agonists, antisense molecules, antibodies, or antagonists. The invention also provides methods for treating disorders associated with **expression** of HMAK.

L7 ANSWER 15 OF 32 MEDLINE . DUPLICATE 3

ACCESSION NUMBER: 1999221639 MEDLINE

DOCUMENT NUMBER: 99221639 PubMed ID: 10205158

TITLE: Presence of a pre-apoptotic complex of pro-caspase-3, Hsp60 and Hsp10 in the **mitochondrial** fraction of jurkat cells.

AUTHOR: Samali A; Cai J; Zhivotovsky B; Jones D P; Orrenius S

CORPORATE SOURCE: Institute of Environmental Medicine, Division of Toxicology, Karolinska Institutet, Box 210, S-171 77, Stockholm, Sweden.. afshin.samali@imm.ki.se

CONTRACT NUMBER: ES09047 (NIEHS)

SOURCE: EMBO JOURNAL, (1999 Apr 15) 18 (8) 2040-8.

Journal code: 8208664. ISSN: 0261-4189.

PUB. COUNTRY: ENGLAND: United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199906  
ENTRY DATE: Entered STN: 19990628  
Last Updated on STN: 19990628  
Entered Medline: 19990611

AB Activation of pro-caspase-3 is a central event in the execution phase of apoptosis and appears to serve as the convergence point of different apoptotic signaling pathways. Recently, **mitochondria** were found to play a central role in apoptosis through release of cytochrome c and activation of caspases. Moreover, a sub-population of pro-caspase-3 has been found to be localized to this organelle. In the present study, we demonstrate that pro-caspase-3 is present in the **mitochondrial** fraction of Jurkat T cells in a complex with the chaperone proteins Hsp60 and Hsp10. Induction of apoptosis with staurosporine led to the activation of **mitochondrial** pro-caspase-3 and its dissociation from the Hsps which were released from **mitochondria**. The release of Hsps occurred simultaneously with the release of other **mitochondrial** intermembrane space proteins including cytochrome c and **adenylate kinase**, prior to a loss of **mitochondrial** transmembrane potential. In in vitro systems, **recombinant** Hsp60 and Hsp10 accelerated the activation of pro-caspase-3 by cytochrome c and dATP in an ATP-dependent manner, consistent with their function as chaperones. This finding suggests that the release of **mitochondrial** Hsps may also accelerate caspase activation in the cytoplasm of intact cells.

L7 ANSWER 16 OF 32 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
ACCESSION NUMBER: 1999-00127 BIOTECHDS

TITLE: **Human mitochondrial adenylate-kinase, HMAK;**  
sense, antisense sequence, antibody, agonist and antagonist used for cancer, neurological and immunological disorder diagnosis and therapy

AUTHOR: Hillman J L; Shah P  
PATENT ASSIGNEE: Incyte-Pharm.  
LOCATION: Palo Alto, CA, USA.  
PATENT INFO: WO 9844124 8 Oct 1998  
APPLICATION INFO: WO 1998-US6249 30 Mar 1998  
PRIORITY INFO: US 1997-829027 31 Mar 1997  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
OTHER SOURCE: WPI: 1998-557119 [47]

AB A purified **mitochondrial adenylate-kinase** (EC-2.7.4.3) with a given protein sequence is claimed. Also claimed is a nucleic acid encoding the kinase, of given nucleotide sequence, and that hybridizes, under stringent conditions, with the given nucleic acid sequence. The claims also cover a nucleic acid complementary to the given sequence, and a DNA probe that constitutes part of that complementary sequence. Also covered are an **expression** vector containing the given nucleic acid sequence, a host cell transformed by that vector, and a means of preparing the **adenylate-kinase** by culturing the transformed cell, and recovering the protein. The claims extend to a composition containing the **adenylate-kinase**, and an antibody, agonist and antagonist of the protein. These are used to treat neurological disorders, cancer and immunological disorders. Also claimed is a means of detecting nucleic acids encoding **mitochondrial adenylate-kinase** in a sample using the DNA probe, and detecting the hybridization complex. The nucleic acids can also be administered for gene therapy. (63pp)

L7 ANSWER 17 OF 32 HCAPLUS COPYRIGHT 2003 ACS

ACCESSION NUMBER: 1998:324881 HCAPLUS

DOCUMENT NUMBER: 129:39786

TITLE: Diabetes-mediating proteins and their therapeutic uses

INVENTOR(S): Mose, Larsen Peter; Fey, Stephen J.; Nerup, Jorn; Karlsen, Allan E.; Bjerre, Christensen Ulla; Pociot, Flemming; Andersen, Henrik U.

PATENT ASSIGNEE(S): Mose Larsen, Peter, Den.; Fey, Stephen J.; Nerup, Jorn; Karlsen, Allan E.; Bjerre Christensen, Ulla; Pociot, Flemming; Andersen, Henrik U.

SOURCE: PCT Int. Appl., 145 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

| PATENT NO.    | KIND   | DATE     | APPLICATION NO. | DATE     |
|---------------|--|----------|-----------------|----------|
| WO 9820124    | A2   | 19980514 | WO 1997-IB1627  | 19971024 |
| WO 9820124    | A3   | 19981008 |                 |          |
| W:            | AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM |          |                 |          |
| RW:           | GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG   |          |                 |          |
| JP 2001500614 | T2   | 20010116 | JP 1998-513441  | 19970916 |
| AU 9854070    | A1   | 19980529 | AU 1998-54070   | 19971024 |
| EP 934409     | A2   | 19990811 | EP 1997-947839  | 19971024 |
| R:            | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI   |          |                 |          |
| JP 2001503860 | T2   | 20010321 | JP 1998-520234  | 19971024 |
| JP 2002504806 | T2   | 20020212 | JP 1998-521182  | 19971024 |
| KR 2000052802 | A  | 20000825 | KR 1999-703621  | 19990424 |

PRIORITY APPLN. INFO.:

|                |    |          |
|----------------|----|----------|
| US 1996-29324P | P  | 19961025 |
| US 1996-30088P | P  | 19961105 |
| US 1996-30186P | P  | 19961105 |
| US 1997-897098 | A2 | 19970718 |
| US 1996-31291P | P  | 19960916 |
| US 1996-29325P | P  | 19961025 |
| WO 1997-IB1114 | W  | 19970916 |
| WO 1997-IB1337 | W  | 19971024 |
| WO 1997-IB1627 | W  | 19971024 |

AB Protective and deleterious diabetes-mediating proteins involved in the development of diabetes or in the prevention of diabetes development are identified by differential **expression** during during development of diabetes relative to **expression** in the absence of diabetes development. These proteins are referred to by their position on 10% IEF or NEPHGE 2-dimensional gels. The purified diabetes-mediating proteins are characterized by mol. wt., isoelec. point, and mass spectroscopic characteristics. Galectin-3 (rat and **human**) and mortalin (mouse and **human**), two of the identified proteins from pancreatic islets, were also sequenced. Transgenic animals **expressing** a diabetes-mediating protein, drug screening methods for identifying a test compd. capable of altering the **expression** of a diabetes-mediating protein, and methods of preventing or ameliorating diabetes by administering a compd. capable of altering the **expression** of a diabetes-mediating protein are also provided..

L7 ANSWER 18 OF 32 MEDLINE DUPLICATE 5  
 ACCESSION NUMBER: 1999033072 MEDLINE  
 DOCUMENT NUMBER: 99033072 PubMed ID: 9813319  
 TITLE: Identification of a novel **adenylate kinase** system in the brain: **cloning** of the fourth **adenylate kinase**.  
 AUTHOR: Yoneda T; Sato M; Maeda M; Takagi H  
 CORPORATE SOURCE: First Department of Anatomy, Osaka City University Medical School, 1-4-3 Asahimachi, Abeno-ku, Osaka-shi, Osaka 545-8585, Japan.  
 SOURCE: BRAIN RESEARCH. MOLECULAR BRAIN RESEARCH, (1998 Nov 20) 62 (2) 187-95.  
 Journal code: 8908640. ISSN: 0169-328X.  
 PUB. COUNTRY: Netherlands  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-D85036; GENBANK-D87809  
 ENTRY MONTH: 199902  
 ENTRY DATE: Entered STN: 19990301  
 Last Updated on STN: 20000303  
 Entered Medline: 19990218

AB We identify a novel subtype of **adenylate kinase**, which is the 4th **adenylate kinase** (AK4), in the vertebrate. AK4 mRNA is **expressed** in the mammalian central nervous system in a region-specific manner from the middle stage of embryogenesis to the adulthood in the rodent. The presence of three isozymes of **adenylate kinase** (AK1, AK2 and AK3) that maintains the homeostasis of adenine and guanine nucleotide composition has been reported in the vertebrate. Obtained mouse AK4 cDNA is 3667 bp in size. The predicted open reading frame consists of 223 amino acid residues. Rat AK4 cDNA is also obtained, and the predicted open reading frame is the same length as that of the mouse. The predicted rat AK4 molecule shows 97.8% homology with mouse AK4. Rat AK4 protein is distinct from rat AK3, 53.8% homologous with rat AK3, although the **adenylate kinase** signature and the **mitochondrial** energy transfer protein signature are found in both sequences. Interestingly, rat AK4 is 89.2% homologous with the **human** AK3 over 223 amino acid residues and rat AK3 is 53.7% homologous with the **human** AK3 indicating that the reported **human** AK3 actually belongs to the AK4 group (therefore, it should be referred to as **human** AK4). Although the sequence of AK4 is most similar to that of AK3 among the AK isozymes, its in vivo **expression** is completely different from AK3; AK4 mRNA is **expressed** in the pyramidal cells in the hippocampus (mainly in the subfield CA3), the granular cells in the cerebellum, nasal neuroepithelium and the liver while AK3 mRNA is **expressed** ubiquitously in the body. It is probable that AK4 acts on the specific mechanism of energy metabolism rather than control of the homeostasis of the ADP pool ubiquitously.  
 Copyright 1998 Elsevier Science B.V.

L7 ANSWER 19 OF 32 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.  
 ACCESSION NUMBER: 97300699 EMBASE  
 DOCUMENT NUMBER: 1997300699  
 TITLE: p32 protein, a splicing factor 2-associated protein, is localized in **mitochondrial** matrix and is functionally important in maintaining oxidative phosphorylation.  
 AUTHOR: Muta T.; Kang D.; Kitajima S.; Fujiwara T.; Hamasaki N.  
 CORPORATE SOURCE: D. Kang, Dept. of Clinical Chem./Lab. Med., Kyushu University Fac. of Medicine, 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-82, Japan

SOURCE: Journal of Biological Chemistry, (1997) 272/39  
(24363-24370).  
Refs: 44  
ISSN: 0021-9258 CODEN: JBCHA3

COUNTRY: United States  
DOCUMENT TYPE: Journal; Article  
FILE SEGMENT: 004 Microbiology  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AB **Human** p32, originally **cloned** as a splicing factor 2-associated protein, has been reported to interact with a variety of molecules including **human** immunodeficiency virus Tat and complement 1q (Clq). p32 protein is supposed to be in the nucleus and on the plasma membrane for the association with **human** immunodeficiency virus Tat and Clq, respectively. None of the interactions, however, is proven to have a physiological role. To investigate the physiological function of p32, we determined the intracellular localization of p32. The fractionation of cells, fluorescent immunocytochemistry, and electron microscopic immunostaining show that p32 is exclusively localized in the **mitochondrial** matrix. We **cloned** a *Saccharomyces cerevisiae* homologue of **human** p32 gene, referred to yeast p30 gene. The yeast p30 protein is also localized in the **mitochondrial** matrix. The disruption of the p30 gene caused the growth retardation of yeast cells in a glycerol medium but not in a glucose medium, i.e. the impairment of the **mitochondrial** ATP synthesis. The growth impairment was restored by the introduction of the **human** p32 cDNA, indicating that p30 is a functional yeast counterpart of **human** p32. Taken together, both p32 and p30 reside in **mitochondrial** matrix and play an important role in maintaining **mitochondrial** oxidative phosphorylation.

L7 ANSWER 20 OF 32 MEDLINE DUPLICATE 6  
ACCESSION NUMBER: 1998088919 MEDLINE  
DOCUMENT NUMBER: 98088919 PubMed ID: 9428643  
TITLE: Intrinsic nucleoside diphosphate kinase-like activity as a novel function of 14-3-3 proteins.  
AUTHOR: Yano M; Mori S; Niwa Y; Inoue M; Kido H  
CORPORATE SOURCE: Division of Enzyme Chemistry, Institute for Enzyme Research, The University of Tokushima, Japan.  
SOURCE: FEBS LETTERS, (1997 Dec 15) 419 (2-3) 244-8.  
Journal code: 0155157. ISSN: 0014-5793.  
PUB. COUNTRY: Netherlands  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199801  
ENTRY DATE: Entered STN: 19980206  
Last Updated on STN: 19980206  
Entered Medline: 19980127

AB 14-3-3 proteins play a role in many cellular functions as molecular chaperone and adapter proteins: they bind to and modulate several proteins involved in cell proliferation and differentiation, and also function ATP-dependently in targeting of precursors to **mitochondria**. We show here that 14-3-3 purified from a **human** lymphoblastoma and also its **recombinant** tau isoform exhibited intrinsic nucleoside diphosphate (NDP) kinase-like activity. 14-3-3 proteins preferentially catalyzed the transfer of the gamma-phosphate group from ATP, dATP or dGTP to all nucleoside diphosphates and this transfer involved acid-labile phosphoenzyme intermediates. They also simultaneously catalyzed the reverse reaction of ATP hydrolysis. These properties of 14-3-3 are similar to those of NDP kinase, but not to those of **adenylate** kinase.

L7 ANSWER 21 OF 32 SCISEARCH COPYRIGHT 2003 THOMSON ISI

ACCESSION NUMBER: 95:187668 SCISEARCH

THE GENUINE ARTICLE: QK489

TITLE: CONTROL OF CELLULAR RESPIRATION IN-VIVO BY  
**MITOCHONDRIAL** OUTER-MEMBRANE AND BY  
CREATINE-KINASE - A NEW SPECULATIVE HYPOTHESIS - POSSIBLE  
INVOLVEMENT OF **MITOCHONDRIAL**-CYTOSKELETON  
INTERACTIONS

AUTHOR: SAKS V A (Reprint); KUZNETSOV A V; KHUCHUA Z A; VASILYEVA  
E V; BELIKOVA J O; KESVATERA T; TIIVEL T

CORPORATE SOURCE: UNIV GRENoble 1, PHYSIOL CELLULAIRE CARDIAQUE LAB, BP 53X,  
F-38041 GRENoble, FRANCE (Reprint); INST CHEM & BIOL PHYS,  
BIOENERGET LAB, TALLINN, ESTONIA; CARDIOL RES CTR,  
BIOENERGET GRP, MOSCOW 121552, RUSSIA

COUNTRY OF AUTHOR: FRANCE; ESTONIA; RUSSIA

SOURCE: JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY, (JAN 1995)  
Vol. 27, No. 1, pp. 625-645.  
ISSN: 0022-2828.

DOCUMENT TYPE: General Review; Journal

FILE SEGMENT: LIFE

LANGUAGE: ENGLISH

REFERENCE COUNT: 119

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB The current problems of regulation of myocardial energy metabolism and oxidative phosphorylation in vivo are considered. With this purpose, retarded diffusion of ADP in cardiomyocytes was studied by analysis of elevated apparent K-m for this substrate in regulation of respiration of saponin-skinned cardiac fibers, as compared to isolated **mitochondria**. Recently published data showing the importance of the outer **mitochondrial** membrane were compared with new experimental results on the proteolysis of skinned fibers and tissue homogenates. In both cases 10 min incubation and 0.125 mg/ml of trypsin resulted in a decrease of apparent K-m for ADP from 297 +/- 35 and 228 +/- 16 to 109 +/- 2 and 36 +/- 16, respectively. Thus, the permeability of the outer **mitochondrial** membrane for ADP may be controlled by some unknown cytoplasmic protein(s), probably related to the cytoskeleton, which are separated from **mitochondria** during their isolation. The extent of **expression** of this protein(s) depends on the energy state and type of muscle. Activation of **mitochondrial** creatine kinase reaction coupled to oxidative phosphorylation overcomes the diffusion difficulties of ADP by amplifying the stimulatory effect of ADP on respiration. It is concluded that both cytoplasmic and **mitochondrial** creatine kinases, **adenylate kinase** and cytoplasmic factor controlling outer membrane permeability may participate in metabolic feedback regulation of respiration in muscle cells.

L7 ANSWER 22 OF 32 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1995:481319 BIOSIS

DOCUMENT NUMBER: PREV199598495619

TITLE: Transfection of a myc gene as a means of generating infinite life span **human** fibroblast strains.

AUTHOR(S): McCormick, J. Justin (1); Kohler, Suzanne K.; Maher, Veronica M.

CORPORATE SOURCE: (1) Carcinogenesis Lab., Fee Hall, Mich. State Univ., East Lansing, MI 48824-1316 USA

SOURCE: Methods in Cell Science, (1995) Vol. 17, No. 2, pp. 141-148.  
ISSN: 1381-5741.

DOCUMENT TYPE: Article

LANGUAGE: English



AB **Human** fibroblasts in culture have never been found to transform spontaneously into immortal cells. In an effort to generate an infinite life span cell strain from foreskin-derived normal diploid fibroblasts, we transfected the cells with a plasmid carrying a v-myc oncogene linked to the neo gene, or with a control vector carrying the neo gene, and selected drug-resistant **clones**. A **clone** that **expressed** the v-myc protein was propagated to the end of its life span, with periodic cryogenic storage of the progeny. The population went into crisis at the same time as cells from the control population and eventually senesced. However, while the cells were senescing, viable-appearing **clones** were noted. The cells of these **clones** continued to multiply, very slowly at first but eventually at a faster rate. Analysis showed that these cells have a diploid karyotype that has remained stable throughout more than 200 population doublings since their sibling cells senesced. Molecular analysis showed that the infinite life span cells are, indeed, derived from the cells used for transfection, and that they continue to **express** the v-myc protein.

L7 ANSWER 23 OF 32 SCISEARCH COPYRIGHT 2003 THOMSON ISI

ACCESSION NUMBER: 94:686537 SCISEARCH

THE GENUINE ARTICLE: PN491

TITLE: PRIMARY AMINO-ACID-SEQUENCE AND STRUCTURE OF **HUMAN** PYRUVATE-CARBOXYLASE

AUTHOR: WEXLER I D (Reprint); DU Y F; LISGARIS M V; MANDAL S K; FREYTAG S O; YANG B S; LIU T C; KWON M; PATEL M S; KERR D S

CORPORATE SOURCE: CASE WESTERN RESERVE UNIV, RAINBOW BABIES & CHILDRENS HOSP, SCH MED, DEPT BIOCHEM, 2047 ABINGTON RD, CLEVELAND, OH, 44106 (Reprint); CASE WESTERN RESERVE UNIV, UNIV HOSP CLEVELAND, SCH MED, DEPT PEDIAT, CLEVELAND, OH, 44106; HENRY FORD HOSP, MOLEC BIOL RES PROGRAM, DETROIT, MI, 48202

COUNTRY OF AUTHOR: USA

SOURCE: BIOCHIMICA ET BIOPHYSICA ACTA-MOLECULAR BASIS OF DISEASE, (21 OCT 1994) Vol. 1227, No. 1-2, pp. 46-52. ISSN: 0925-4439.

DOCUMENT TYPE: Article; Journal

FILE SEGMENT: LIFE

LANGUAGE: ENGLISH

REFERENCE COUNT: 43

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB Pyruvate carboxylase (PC) (pyruvate:carbon dioxide ligase (ADP-forming), EC 6.4.1.1.), a nuclear-encoded **mitochondrial** enzyme, catalyzes the conversion of pyruvate to oxaloacetate. We have isolated and characterized cDNAs spanning the entire coding region of **human** PC. The sequence of **human** PC has an open reading frame of 3537 nucleotides which encodes for a polypeptide with a length of 1178 amino acids. The identity of the cDNA as PC is confirmed by comparison to PC cDNAs of other species and sequenced peptide fragments of mammalian PC. The M(r) of the full length precursor protein is 129 576 and that of the mature apoprotein is 127 370. RNA blot analysis from a variety of **human** tissues demonstrates that the highest level of PC mRNA is found in liver corresponding to this tissue's high level of PC activity. Based on homology with other biotin-containing proteins, the ATP, pyruvate, and biotin-binding sites can be identified. One of two patients with documented PC deficiency was found to be missing PC mRNA, further confirming the identity of this cDNA.

L7 ANSWER 24 OF 32 SCISEARCH COPYRIGHT 2003 THOMSON ISI

ACCESSION NUMBER: 92:709675 SCISEARCH

THE GENUINE ARTICLE: KB012

TITLE: VIRAL THYMIDINE KINASES AND THEIR RELATIVES

AUTHOR: GENTRY G A (Reprint)  
CORPORATE SOURCE: UNIV MISSISSIPPI, MED CTR, DEPT MICROBIOL, JACKSON, MS, 39216 (Reprint)  
COUNTRY OF AUTHOR: USA  
SOURCE: PHARMACOLOGY & THERAPEUTICS, (1992) Vol. 54, No. 3, pp. 319-355.  
ISSN: 0163-7258.  
DOCUMENT TYPE: General Review; Journal  
FILE SEGMENT: LIFE  
LANGUAGE: ENGLISH  
REFERENCE COUNT: 200

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB Thymidine kinases were described for cellular life long before it was shown that they could also be encoded by viruses, but the viral thymidine kinase genes were the first to be sequenced. These enzymes have been extraordinarily useful to the researcher, serving first to help label DNA, then to get thymidine analogs incorporated into DNA for therapeutic and other purposes and more recently to move genes from one genome to another. Knowledge of the nucleotide and amino acid sequences of these enzymes has allowed some deductions about their possible three-dimensional structure, as well as the location on the polypeptide of various functions; it has also allowed their classification into two main groups: the herpesviral thymidine/eukaryotic deoxycytidine kinases and the poxviral and cellular thymidine kinases; the relationships of the **mitochondrial** enzyme are still not clear.

L7 ANSWER 25 OF 32 MEDLINE

ACCESSION NUMBER: 90363911 MEDLINE  
DOCUMENT NUMBER: 90363911 PubMed ID: 2168054  
TITLE: Gene structures of three vertebrate **adenylate kinase** isozymes.  
AUTHOR: Nakazawa A; Yamada M; Tanaka H; Shahjahan M; Tanabe T  
CORPORATE SOURCE: Department of Biochemistry, Yamaguchi University School of Medicine, Japan.  
SOURCE: PROGRESS IN CLINICAL AND BIOLOGICAL RESEARCH, (1990) 344 495-514.  
Journal code: 7605701. ISSN: 0361-7742.  
PUB. COUNTRY: United States  
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199010  
ENTRY DATE: Entered STN: 19901109  
Last Updated on STN: 19901109  
Entered Medline: 19901003

AB **Adenylate kinase** is an ubiquitous enzyme which contributes to homeostasis of adenine nucleotide composition in the cell. In vertebrates, three isozymes (AK1, AK2, and AK3) are characterized which have distinct distribution in tissues as well as subcellular compartments. The genetic backgrounds of these **adenylate kinase** isozymes were analyzed. cDNA **clones** for AK1 were isolated from poly(A)+RNA of chicken skeletal muscle. The results of mRNA analysis in various tissues using the AK1 cDNA indicated that the AK1 gene **expression** is regulated both tissue-specifically and developmentally at the transcriptional level. The AK1 gene was **cloned** from chicken and **human** DNA and characterized. Both genes were split into seven exons. The intron positions in both genes coincided. cDNA **clones** for AK2 isolated from bovine liver poly(A)+RNA contained two types. One type (AK2A) encoded the same amino acid sequence as that reported for bovine heart AK2. The other type (AK2B) encoded the same sequence as AK2 except for the COOH terminus. The mRNA species corresponding to the two cDNA **clones** were

identified in bovine liver and heart.. Both the cDNA sequences were found to direct the active **adenylate kinase** synthesis in *E. coli*. The AK2 gene was **cloned** and characterized. It consisted of seven exons and six introns. From genomic structure analysis, the two cDNA species were shown to be derived from a single gene by the alternative splicing mechanism. Three types of cDNA **clones** for AK3 were isolated from bovine liver poly(A)+RNA, which contained the common AK3-coding region and different 3' portions. No NH2-terminal presequence of **mitochondrial** targeting was identified in AK3 from the sequencing and **expression** analyses of the cDNA. Upon **expression** of the cDNA sequence in *E. coli*, AK3 protein was recovered in the periplasmic space of the bacteria, indicating that AK3 without presequence was exported through the inner bacterial membrane as it is imported through the **mitochondrial** membranes. Internal targeting signals may be responsible for the translocation process. The AK3 gene was **cloned** and partially characterized. It is split into at least five exons. The comparisons of amino acid sequences and genomic structure of three isozymes revealed that a segment corresponding to either exon 5 of the AK2 gene or a part of exon 3 of the AK3 gene is missing in the AK1 gene. Phylogenetic analysis suggested that AK1, a shorter molecule, would have been separated from a longer molecule very early in evolution of **adenylate kinase**. (ABSTRACT TRUNCATED AT 400 WORDS)

L7 ANSWER 26 OF 32 MEDLINE  
 ACCESSION NUMBER: 90037053 MEDLINE  
 DOCUMENT NUMBER: 90037053 PubMed ID: 2478555  
 TITLE: **Cloning** and characterization of cDNA for **mitochondrial** GTP:AMP phosphotransferase of bovine liver.  
 AUTHOR: Yamada M; Shahjahan M; Tanabe T; Kishi F; Nakazawa A  
 CORPORATE SOURCE: Department of Biochemistry, Yamaguchi University School of Medicine, Japan.  
 SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1989 Nov 15) 264 (32) 19192-9.  
 Journal code: 2985121R. ISSN: 0021-9258.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 OTHER SOURCE: GENBANK-M25757  
 ENTRY MONTH: 198912  
 ENTRY DATE: Entered STN: 19900328  
 Last Updated on STN: 19960129  
 Entered Medline: 19891215

AB Three different types of cDNA **clones** for **mitochondrial** GTP:AMP phosphotransferase (AK3) were isolated from a cDNA library of bovine liver poly(A)+ RNA. Nucleotide sequencing revealed that each of these **clones** consisted of a common 5'-untranslated region, a common AK3-coding sequence and a 3'-untranslated region with different sizes. By Northern blot analysis, three species of AK3 mRNA apparently corresponding to the isolated cDNA **clones** were detected, which would be a result of varying terminations and polyadenylations of the primary transcript. From comparison of the size of the product synthesized *in vitro* from the message directed by the isolated cDNA with that of the purified AK3 protein, AK3 appeared to have no cleavable NH2-terminal sequence as found in other **mitochondrial** proteins. The AK3 cDNA was **expressed** in *Escherichia coli*, which resulted in complementation of an **adenylate kinase** mutation of *E. coli*. The AK3 product was exported to the periplasmic space through the bacterial inner membrane. The possible involvement of the NH2-terminal sequence of the protein in targeting to the

mitochondrial matrix was discussed.

L7 ANSWER 27 OF 32 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.DUPLICATE 7  
ACCESSION NUMBER: 83053333 EMBASE  
DOCUMENT NUMBER: 1983053333  
TITLE: Adenosine triphosphate-adenosine-5'-monophosphate  
phosphotransferase from normal **human** liver  
**mitochondria**. Isolation, chemical properties, and  
immunochemical comparison with Duchenne dystrophic serum  
aberrant **adenylate kinase**.  
AUTHOR: Hamada M.; Sumida M.; Okuda H.; et al.  
CORPORATE SOURCE: Dep. Hyg., Ehime Univ. Sch. Med., Shigenobu cho, Onsen gun,  
Ehime 791-02, Japan  
SOURCE: Journal of Biological Chemistry, (1982) 257/21  
(13120-13128).  
CODEN: JBCHA3  
COUNTRY: United States  
DOCUMENT TYPE: Journal  
FILE SEGMENT: 029 Clinical Biochemistry  
008 Neurology and Neurosurgery  
LANGUAGE: English

AB **Adenylate kinase** has been purified approximately  
1360-fold to a final specific activity of 280 .mu.mol of ATP formed  
min-1xmg-1 of protein at 30.degree.C from normal **human** liver  
**mitochondria**. The purity of the final preparation was evaluated by  
studies with polyacrylamide gel electrophoresis and sodium dodecyl  
sulfate-polyacrylamide gel electrophoresis and by sedimentation studies.  
The purified enzyme catalyzes transphosphorylation reactions between  
adenosine triphosphate (ATP) and adenosine monophosphate (AMP). ATP and  
adenosine-5'-thiophosphate, ATP and adenosine monophosphate-3'-  
pyrophosphate, adenosine-s'-(3-thio)triphosphate and AMP. The nearly  
constant ratios of these activities throughout the purification scheme  
suggest that all are catalyzed by the same enzyme. The purified enzyme has  
a molecular weight of 25,200 by sedimentation equilibrium with the use of  
a partial specific volume of 0.73 mlxg-1 calculated from amino acid  
analysis. This purified enzyme was also found to be a single polypeptide  
with a molecular weight of 26,500 by sodium dodecyl sulfate-polyacrylamide  
gel electrophoresis. From amino acid analysis, a calculated minimum  
molecular weight of 26,349 was obtained. Initial velocity studies revealed  
a narrow specificity for adenine nucleotides. The Kd' values for MgATP2-  
and MgATP2-.gamma.S1 were 0.12 and 0.57 .mu.M with Vmax.forward values of  
1.04 (.+-0.04)x103 and 7.02x102 .mu.mol x min-1 x mg-1, respectively. For  
the monophosphate acceptor, Kd' values of 0.56 and 186 .mu.M were measured  
for 5'-AMP2- and AMP2-.alpha.S, respectively. The Kd' for MgADP1- and  
ADP3- were 0.53 and 0.17 .mu.M with a Vmax.reverse of 6.40(.+-0.03)x102  
.mu.molxmin-1xmg-1 of protein. The steady state kinetics, at pH 7.4,  
30.degree.C, and essentially fixed .DELTA./2 of 0.16-0.18, of this enzyme  
seem to be adequately **expressed** by a random quasi-equilibrium  
type of mechanism with a rate-limiting step largely at the interconversion  
of the ternary complexes, as shown in rabbit muscle, calf muscle, and calf  
liver **adenylate kinase**. It would appear that normal  
**human** liver **mitochondrial adenylate**  
**kinase** largely favors the forward reaction (ADP formation). A  
specific anti-liver enzyme antibody obtained from rabbit serum inhibited  
the purified liver **mitochondrial** enzyme activity, but not the  
purified **human** muscle enzyme, nor the aberrant **adenylate**  
**kinase** from Duchenne dystrophic serum.

L7 ANSWER 28 OF 32 MEDLINE DUPLICATE 8  
ACCESSION NUMBER: 82003493 MEDLINE  
DOCUMENT NUMBER: 82003493 PubMed ID: 6944169  
TITLE: Characterization of the Philadelphia chromosome by gene

mapping.  
 AUTHOR: Geurts van Kessel A H; ten Brinke H; Boere W A; den Boer W C; de Groot P G; Hagemeijer A; Meera Khan P; Pearson P L  
 SOURCE: CYTOGENETICS AND CELL GENETICS, (1981) 30 (2) 83-91.  
 Journal code: 0367735. ISSN: 0301-0171.  
 PUB. COUNTRY: Switzerland  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 198111  
 ENTRY DATE: Entered STN: 19900316  
 Last Updated on STN: 19900316  
 Entered Medline: 19811122

AB Chinese hamster X **human** and mouse X **human** somatic cell hybrid lines were obtained using circulating leucocytes from six chronic myeloid leukemia patients. All six patients carried the Ph1 translocation, t(9q+;22q-), characteristic of chronic myeloid leukemia, in their dividing immature granulocytes. Analysis of independent hybrid **clones** yielded the following results: 1. The chromosome 9 markers, soluble aconitase and **adenylate kinase-1**, segregated with the 9q+ derivative. The latter marker has previously been localized to 9q34. 2. The chromosome 22 markers, **mitochondrial** aconitase, N-acetyl-alpha-D-galactosaminidase, and arylsulfatase-A, also segregated with the 9q+ derivative. **Mitochondrial** aconitase has recently been assigned to 22q11 leads to 22q13. No evidence was obtained either for reciprocity of the translocation or for variations in breakpoints in different patients. The results reported in this paper provisionally assign the gene for **mitochondrial** aconitase to a region distal to the breakpoint in 22q11.

L7 ANSWER 29 OF 32 LIFESCI COPYRIGHT 2003 CSA

ACCESSION NUMBER: 81:24127 LIFESCI  
 TITLE: Characterization of the Philadelphia Chromosome by Gene Mapping.  
 AUTHOR: Van Kessel, A.H.M.G.; Ten Brinke, H.; Boere, W.A.M.; Den Boer, W.C.; De Groot, P.G.; Hagemeijer, A.; Meera Khan, P.; Pearson, P.L.  
 CORPORATE SOURCE: Dept. Cell Biol. Genet., Erasmus Univ., P.O. Box 1738, 3000 DR Rotterdam, Netherland  
 SOURCE: CYTOGENET. CELL GENET., (1981) vol. 30, no. 2, pp. 83-91.  
 DOCUMENT TYPE: Journal  
 FILE SEGMENT: G  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB Chinese hamster x **human** and mouse x **human** somatic cell hybrid lines were obtained using circulating leucocytes from six chronic myeloid leukemia patients. All six patients carried the Ph super(1) translocation, t(9q+;22q-), characteristic of chronic myeloid leukemia, in their dividing immature granulocytes. Analysis of independent hybrid **clones** yielded the following results: 1. The chromosome 9 markers, soluble aconitase and **adenylate kinase-1**, segregated with the 9q+ derivative. The latter marker has previously been localized to 9q34. 2. The chromosome 22 markers, **mitochondrial** aconitase, N-acetyl- alpha -D-galactosaminidase, and arylsulfatase-A, also segregated with the 9q+ derivative. **Mitochondrial** aconitase has recently been assigned to 22q11 arrow right 22q13. No evidence was obtained either for reciprocity of the translocation or for variations in breakpoints in different patients.

L7 ANSWER 30 OF 32 MEDLINE

ACCESSION NUMBER: 79194246 MEDLINE  
 DOCUMENT NUMBER: 79194246 PubMed ID: 36399

TITLE: Cytosolic phosphorylation potential.  
 AUTHOR: Veech R L; Lawson J W; Cornell N W; Krebs H A  
 SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1979 Jul 25) 254 (14) 6538-47.  
 Journal code: 2985121R. ISSN: 0021-9258.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 197909  
 ENTRY DATE: Entered STN: 19900315  
 Last Updated on STN: 19990129  
 Entered Medline: 19790901

AB The tissue contents of the reactants of the myokinase (EC 2.7.4.3) and the combined glyceraldehyde-3-phosphate dehydrogenase (EC 1.1.1.29)-3-phosphoglycerate kinase (EC 2.7.2.3) reactions were measured in rapidly inactivated samples of **human** blood and rat brain, muscle, and liver. The tissue contents of the reactants of the creatine kinase (EC 2.7.3.2) reaction were measured in rat brain and muscle. In vitro the value of the **expression**:  $KG+G = \frac{[\sigma_3PG]}{[\sigma_{ATP}]}$ .  $\frac{[\sigma_{lactate}]}{KLDH} = \frac{[\sigma_{HAP}]}{22} \cdot \frac{[\sigma_{ADP}][\sigma_{Pi}]}{[\sigma_{RUVATE}]}$  (1) was found to be  $0.725 \times 10(7) \text{ M}^{-1}$  at  $I = 0.25$ ,  $T = 38$  degrees C, and free  $[Mg^{2+}] = 0.15 \text{ mM}$  and the value measured in vivo in red cell was  $0.699 \times 10(7) \text{ M}^{-1}$ . The value of the **expression**  $KMYK = \frac{[\sigma_{ATP}][\sigma_{AMP}]}{[\sigma_{ADP}]}$  measured under the above conditions and at pH 7.2 was found to be 0.744 while the value found in red cell was  $0.784 \pm 0.037$ . These reactions, therefore, appear to be in a state of near-equilibrium in the red cell and the measured tissue contents of ATP and ADP, which are common reactants in both reactions, approximate closely the activity of these reactants in vivo. In brain and muscle, the value of  $KG + G/KLDH$  calculated from the measured tissue contents of the reactants was a factor of 20 or more lower than that expected at equilibrium as was the measured value of the **expression**:  $KCK = \frac{[\sigma_{ATP}][\sigma_{creatine}]}{[\sigma_{ADP}][\sigma_{creatine-P}][H^+]}$  (2) Substitution of calculated free  $[\sigma_{ADP}]$  values in the **expression** of  $KG + G/KLDH$  gave values of  $0.83 \pm 0.19 \times 10(7) \text{ M}^{-1}$  for brain and muscle, respectively, which agreed well with the value of  $1.65 \times 10(7) \text{ M}^{-1}$  measured in vitro at  $I = 0.25$ , free  $[Mg^{2+}] = 1 \text{ mM}$ ,  $T = 38$  degrees C. This agreement between two highly active enzyme systems in the same compartment is taken as evidence of the existence of near-equilibrium in both these systems and suggests that free cytosolic  $[\sigma_{ADP}]$  is probably 20-fold lower than measured cell ADP content in **mitochondrial**-containing tissues.

L7 ANSWER 31 OF 32 MEDLINE DUPLICATE 9  
 ACCESSION NUMBER: 77221531 MEDLINE  
 DOCUMENT NUMBER: 77221531 PubMed ID: 195572  
 TITLE: **Adenylate kinase 2, a mitochondrial enzyme.**  
 AUTHOR: Bruns G A; Regina V M  
 SOURCE: BIOCHEMICAL GENETICS, (1977 Jun) 15 (5-6) 477-86.  
 Journal code: 0126611. ISSN: 0006-2928.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 197709  
 ENTRY DATE: Entered STN: 19900314  
 Last Updated on STN: 19970203  
 Entered Medline: 19770902

AB The subcellular compartmentalization of the isoenzymes of ATP:AMP phosphotransferase (**adenylate kinase**) was analyzed in

HeLa cells, RAG cells, and RAG-human hybrids that express human AK-2. In HeLa cells and in the hybrids, human AK-2 was present in a mitochemical fraction prepared from cell extracts and in mitochondria purified by density gradient centrifugation. Human AK-1 was, as expected, distributed in the soluble cytoplasmic fraction of the cells. The rodent isozymes which are homologous to human AK-1 and AK-2 have been determined.

L7 ANSWER 32 OF 32 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 1977:15881 BIOSIS

DOCUMENT NUMBER: BR13:15881

TITLE: ASSIGNMENT OF HUMAN GENES BETA GLUCURONIDASE TO CHROMOSOME 7 ADENYLATE KINASE 1 TO 9 A 2ND ENZYME WITH ENOLASE ACTIVITY TO 12 AND MITOCHONDRIAL ISO CITRATE DEHYDROGENASE TO 15.

AUTHOR(S): GRZESCHIK K-H

SOURCE: Cytogenet. Cell Genet., (1976) 16 (1-5), 142-148. CODEN: CGCGBR. ISSN: 0301-0171.

FILE SEGMENT: BR; OLD

LANGUAGE: Unavailable

=> s "HMAK"

L8 8 "HMAK"

=> dup rem l8

PROCESSING COMPLETED FOR L8

L9 3 DUP REM L8 (5 DUPLICATES REMOVED)

=> d 1-3 ibib ab

L9 ANSWER 1 OF 3 MEDLINE DUPLICATE 1

ACCESSION NUMBER: 2002485451 MEDLINE

DOCUMENT NUMBER: 22217966 PubMed ID: 12084720

TITLE: Identification of human male germ cell-associated kinase, a kinase transcriptionally activated by androgen in prostate cancer cells.

AUTHOR: Xia Liang; Robinson Dan; Ma Ai-Hong; Chen Hua-Chien; Wu Frederick; Qiu Yun; Kung Hsing-Jien

CORPORATE SOURCE: Department of Biological Chemistry, School of Medicine, University of California, Davis, California 95616, USA.

CONTRACT NUMBER: CA39207 (NCI)

CA57179 (NCI)

CA82073 (NCI)

DK52659 (NIDDK)

SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (2002 Sep 20) 277 (38) 35422-33.

Journal code: 2985121R. ISSN: 0021-9258.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

OTHER SOURCE: GENBANK-AF505623

ENTRY MONTH: 200210

ENTRY DATE: Entered STN: 20020926

Last Updated on STN: 20030105

Entered Medline: 20021024

AB Androgen is involved in both normal development and malignant transformation of prostate cells. The signal transduction pathways associated with these processes are not well understood. Using a novel kinase display approach, we have identified a protein kinase, human male germ cell-associated kinase (hMAK), which is transcriptionally

induced by the androgenic hormone 5 $\alpha$ -dihydrotestosterone (DHT). The kinetics of induction is rapid and dose-dependent, and the induction is not blocked by cycloheximide treatment. Real time reverse transcription-PCR studies demonstrated a 9-fold induction of **hMAK** by 10 nM DHT at 24 h post-stimulation. The expression levels of **hMAK** in prostate cancer cell lines are in general higher than those of normal prostate epithelial cells. A reverse transcription-PCR product encompassing the entire **hMAK** open reading frame was isolated. The results from sequencing analysis showed that the **hMAK** protein is 623 amino acids in length and contains a kinase catalytic domain at its N terminus, followed by a proline/glutamine-rich domain. The catalytic domain of this kinase contains sequence motifs related to both the cyclin-dependent kinase and the mitogen-activated protein kinase families. When expressed in COS1 cells, **hMAK** is kinase-active as demonstrated by autophosphorylation and phosphorylation of exogenous substrate and is localized in the nucleus. A 3.7-kilobase pair promoter of the **hMAK** locus was isolated from a human genomic DNA bacterial artificial chromosome clone and was shown to be activated by DHT. This activation can be blocked by an anti-androgen drug bicalutamide (Casodex), implicating the involvement of androgen receptor in this process. Taken together, these data suggest that **hMAK** is a protein kinase targeted by androgen that may participate in androgen-mediated signaling in prostate cancer cells.

L9 ANSWER 2 OF 3 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.  
 ACCESSION NUMBER: 2000:290852 BIOSIS  
 DOCUMENT NUMBER: PREV200000290852  
 TITLE: Mitochondrial adenylate kinase.  
 AUTHOR(S): Hillman, Jennifer L.; Shah, Purvi  
 ASSIGNEE: Incyte Pharmaceuticals, Inc.  
 PATENT INFORMATION: US 6001624 December 14, 1999  
 SOURCE: Official Gazette of the United States Patent and Trademark Office Patents, (Dec. 14, 1999) Vol. 1229, No. 2, pp. No pagination. e-file.  
 ISSN: 0098-1133.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English

AB The present invention provides a human mitochondrial adenylate kinase (**hMAK**) and polynucleotides which encode **hMAK**. The invention also provides expression vectors, host cells, agonists, antisense molecules, antibodies, or antagonists. The invention also provides methods for treating disorders associated with expression of **hMAK**.

L9 ANSWER 3 OF 3 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
 ACCESSION NUMBER: 1999-00127 BIOTECHDS  
 TITLE: Human mitochondrial adenylate-kinase, **hMAK**;  
 sense, antisense sequence, antibody, agonist and antagonist used for cancer, neurological and immunological disorder diagnosis and therapy  
 AUTHOR: Hillman J L; Shah P  
 PATENT ASSIGNEE: Incyte-Pharm.  
 LOCATION: Palo Alto, CA, USA.  
 PATENT INFO: WO 9844124 8 Oct 1998  
 APPLICATION INFO: WO 1998-US6249 30 Mar 1998  
 PRIORITY INFO: US 1997-829027 31 Mar 1997  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 OTHER SOURCE: WPI: 1998-557119 [47]

AB A purified mitochondrial adenylate-kinase (EC-2.7.4.3) with a given protein sequence is claimed. Also claimed is a nucleic acid encoding the kinase, of given nucleotide sequence, and that hybridizes, under



stringent conditions, with the given nucleic acid sequence. The claims also cover a nucleic acid complementary to the given sequence, and a DNA probe that constitutes part of that complementary sequence. Also covered are an expression vector containing the given nucleic acid sequence, a host cell transformed by that vector, and a means of preparing the adenylate-kinase by culturing the transformed cell, and recovering the protein. The claims extend to a composition containing the adenylate-kinase, and an antibody, agonist and antagonist of the protein. These are used to treat neurological disorders, cancer and immunological disorders. Also claimed is a means of detecting nucleic acids encoding mitochondrial adenylate-kinase in a sample using the DNA probe, and detecting the hybridization complex. The nucleic acids can also be administered for gene therapy. (63pp)

=> e hillman j l/au

|     |         |                      |
|-----|---------|----------------------|
| E1  | 77      | HILLMAN J J/AU       |
| E2  | 5       | HILLMAN J K/AU       |
| E3  | 454 --> | HILLMAN J L/AU       |
| E4  | 1       | HILLMAN J M/AU       |
| E5  | 4       | HILLMAN J M L/AU     |
| E6  | 2       | HILLMAN J O/AU       |
| E7  | 5       | HILLMAN J P/AU       |
| E8  | 242     | HILLMAN J R/AU       |
| E9  | 2       | HILLMAN J RICHARD/AU |
| E10 | 109     | HILLMAN J S/AU       |
| E11 | 21      | HILLMAN J T/AU       |
| E12 | 10      | HILLMAN J V/AU       |

=> s e3

L10 454 "HILLMAN J L"/AU

=> d his

(FILE 'HOME' ENTERED AT 12:28:11 ON 06 MAY 2003)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:28:34 ON 06 MAY 2003

|     |  |
|-----|--|
| L1  | 10975 S ADENYLATE KINASE?                  |
| L2  | 2615 S HUMAN AND L1                        |
| L3  | 613747 S MITOCHONDR?                       |
| L4  | 255 S L2 AND L3                            |
| L5  | 5929362 S CLON? OR EXPRESS? OR RECOMBINANT |
| L6  | 59 S L4 AND L5                             |
| L7  | 32 DUP REM L6 (27 DUPLICATES REMOVED)      |
| L8  | 8 S "HMAK"                                 |
| L9  | 3 DUP REM L8 (5 DUPLICATES REMOVED)        |
|     | E HILLMAN J L/AU                           |
| L10 | 454 S E3                                   |

=> s l4 and l10

L11 1 L4 AND L10

=> d all

|     |  |
|-----|--|
| L11 | ANSWER 1 OF 1 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI   |
| AN  | 1999-00127 BIOTECHDS   |
| TI  | <b>Human mitochondrial adenylate-kinase, HMAK;</b><br>sense, antisense sequence, antibody, agonist and antagonist used for cancer, neurological and immunological disorder diagnosis and therapy |
| AU  | Hillman J L; Shah P  |

PA Incyte-Pharm.  
 LO Palo Alto, CA, USA.  
 PI WO 9844124 8 Oct 1998  
 AI WO 1998-US6249 30 Mar 1998  
 PRAI US 1997-829027 31 Mar 1997  
 DT Patent  
 LA English  
 OS WPI: 1998-557119 [47]  
 AB A purified **mitochondrial adenylate-kinase** (EC-2.7.4.3) with a given protein sequence is claimed. Also claimed is a nucleic acid encoding the kinase, of given nucleotide sequence, and that hybridizes, under stringent conditions, with the given nucleic acid sequence. The claims also cover a nucleic acid complementary to the given sequence, and a DNA probe that constitutes part of that complementary sequence. Also covered are an expression vector containing the given nucleic acid sequence, a host cell transformed by that vector, and a means of preparing the **adenylate-kinase** by culturing the transformed cell, and recovering the protein. The claims extend to a composition containing the **adenylate-kinase**, and an antibody, agonist and antagonist of the protein. These are used to treat neurological disorders, cancer and immunological disorders. Also claimed is a means of detecting nucleic acids encoding **mitochondrial adenylate-kinase** in a sample using the DNA probe, and detecting the hybridization complex. The nucleic acids can also be administered for gene therapy. (63pp)  
 CC D PHARMACEUTICALS; D3 Peptides and Proteins; D PHARMACEUTICALS; D7 Clinical Genetic Techniques; D PHARMACEUTICALS; D6 Antibodies; A GENETIC ENGINEERING AND FERMENTATION; A1 Nucleic Acid Technology  
 CT **HUMAN MITOCHONDRIA RECOMBINANT ADENYLATE-KINASE** PREP., VECTOR EXPRESSION IN HOST CELL, DNA PROBE, ANTIBODY, AGONIST, ANTAGONIST, SENSE, ANTISENSE SEQUENCE, APPL. CANCER, NEUROLOGICAL DISORDER, IMMUNOLOGICAL DISORDER THERAPY, GENE THERAPY, DIAGNOSIS ANIMAL MAMMAL ENZYME EC-2.7.4.3 DNA SEQUENCE PROTEIN SEQUENCE GENE TRANSFER HYBRIDIZATION CLONING TUMOR (VOL.18, NO.1)

=> e shah p/au

|     |          |                     |
|-----|----------|---------------------|
| E1  | 1        | SHAH OMAR JAMEEL/AU |
| E2  | 2        | SHAH OMAR JAVED/AU  |
| E3  | 1409 --> | SHAH P/AU           |
| E4  | 131      | SHAH P A/AU         |
| E5  | 115      | SHAH P B/AU         |
| E6  | 214      | SHAH P C/AU         |
| E7  | 51       | SHAH P D/AU         |
| E8  | 64       | SHAH P G/AU         |
| E9  | 3        | SHAH P G S/AU       |
| E10 | 57       | SHAH P H/AU         |
| E11 | 5        | SHAH P I/AU         |
| E12 | 153      | SHAH P J/AU         |

=> s e3

L12 1409 "SHAH P"/AU

=> df his

DF IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.  
 For a list of commands available to you in the current file, enter "HELP COMMANDS" at an arrow prompt (=>).

=> d his

(FILE 'HOME' ENTERED AT 12:28:11 ON 06 MAY 2003)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS, LIFESCI' ENTERED AT 12:28:34 ON 06 MAY 2003

L1 10975 S ADENYLATE KINASE?  
L2 2615 S HUMAN AND L1  
L3 613747 S MITOCHONDR?  
L4 255 S L2 AND L3  
L5 5929362 S CLON? OR EXPRESS? OR RECOMBINANT  
L6 59 S L4 AND L5  
L7 32 DUP REM L6 (27 DUPLICATES REMOVED)  
L8 8 S "HMAK"  
L9 3 DUP REM L8 (5 DUPLICATES REMOVED)  
E HILLMAN J L/AU  
L10 454 S E3  
L11 1 S L4 AND L10  
E SHAH P/AU  
L12 1409 S E3

=> s l4 and l12

L13 1 L4 AND L12

=> d ibib ab

L13 ANSWER 1 OF 1 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI  
ACCESSION NUMBER: 1999-00127 BIOTECHDS

TITLE: **Human mitochondrial adenylate-kinase, HMAK;**  
sense, antisense sequence, antibody, agonist and  
antagonist used for cancer, neurological and immunological  
disorder diagnosis and therapy

AUTHOR: Hillman J L; Shah P  
PATENT ASSIGNEE: Incyte-Pharm.  
LOCATION: Palo Alto, CA, USA.  
PATENT INFO: WO 9844124 8 Oct 1998  
APPLICATION INFO: WO 1998-US6249 30 Mar 1998  
PRIORITY INFO: US 1997-829027 31 Mar 1997  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
OTHER SOURCE: WPI: 1998-557119 [47]

AB A purified **mitochondrial adenylate-kinase** (EC-2.7.4.3) with a given protein sequence is claimed. Also claimed is a nucleic acid encoding the kinase, of given nucleotide sequence, and that hybridizes, under stringent conditions, with the given nucleic acid sequence. The claims also cover a nucleic acid complementary to the given sequence, and a DNA probe that constitutes part of that complementary sequence. Also covered are an expression vector containing the given nucleic acid sequence, a host cell transformed by that vector, and a means of preparing the **adenylate-kinase** by culturing the transformed cell, and recovering the protein. The claims extend to a composition containing the **adenylate-kinase**, and an antibody, agonist and antagonist of the protein. These are used to treat neurological disorders, cancer and immunological disorders. Also claimed is a means of detecting nucleic acids encoding **mitochondrial adenylate-kinase** in a sample using the DNA probe, and detecting the hybridization complex. The nucleic acids can also be administered for gene therapy. (63pp)

=> d his

(FILE 'HOME' ENTERED AT 12:28:11 ON 06 MAY 2003)

FILE 'MEDLINE, EMBASE, BIOSIS, BIOTECHDS, SCISEARCH, HCAPLUS, NTIS,  
LIFESCI' ENTERED AT 12:28:34 ON 06 MAY 2003

L1 10975 S ADENYLATE KINASE?  
L2 2615 S HUMAN AND L1  
L3 613747 S MITOCHONDR?  
L4 255 S L2 AND L3  
L5 5929362 S CLON? OR EXPRESS? OR RECOMBINANT  
L6 59 S L4 AND L5  
L7 32 DUP REM L6 (27 DUPLICATES REMOVED)  
L8 8 S "HMAK"  
L9 3 DUP REM L8 (5 DUPLICATES REMOVED)  
E HILLMAN J L/AU  
L10 454 S E3  
L11 1 S L4 AND L10  
E SHAH P/AU  
L12 1409 S E3  
L13 1 S L4 AND L12